

Oxalate impacts macrophage metabolism and immune response to uropathogenic *E. coli* infection

Parveen Kumar, PhD¹, Vikram Saini, PhD², and Tanecia Mitchell, PhD¹

¹Department of Urology, University of Alabama at Birmingham, Birmingham, AL, USA, ²All India Institute of Medical Sciences, New Delhi, India

Introduction & Objective

INTRODUCTION

- Urinary stones are suggested to be associated with oxalate and bacterial infection.
- Escherichia coli* (*E. coli*) has been isolated from human urine and urinary stones composed of oxalate.¹⁻² It has also been shown to promote calcium oxalate (CaOx) crystallization.³
- Stone formers have a high incidence of urinary tract infection (UTI) and increased inflammation.⁴⁻⁵
- Oxalate and crystals have been shown to induce inflammation (i.e. Monocyte Chemoattractant Protein-1 chemokine release) in renal epithelial cells⁶⁻⁷.
- The mechanisms leading to UTI in kidney stone patients are not well defined.
- We have previously determined that 1) monocyte cellular energetics is reduced in CaOx kidney stone formers, and 2) oxalate alters redox homeostasis and cellular energetics in monocytes (Figure 1).





Figure 1: Cellular energetic profiles in (A) monocytes isolated from healthy subjects (HS; black circle) and CaOx SF (white circle) and (B) monocytes from HS (black circle) treated with sodium oxalate. Results are presented as mean $\pm$ se; n=5-6 replicates per group.

OBJECTIVE

—To determine the impact of oxalate on macrophage metabolism, redox status, and anti-bacterial response to uropathogenic *Escherichia coli* (*E. coli*) infection.

Methods



Methods

- Cell Viability & Proliferation:** CCK8 and LDH assays were performed to assess macrophage proliferation and viability, respectively.
- Mitochondrial Membrane Potential:** was measured using TMRE, a fluorescence dye.
- Oxidative Stress:** Mitochondrial superoxide was measured using a fluorescence mitochondrial superoxide dye and total ROS was assessed using DCF staining.
- Cellular Energetics:** A Seahorse XF Analyzer was used to assess metabolism.
- Infection assay:** The colony forming unit (CFU) count was determined by counting the amount of bacteria present in cell lysates following treatment.

Results




Figure 2: The effect of oxalate and CFT037 on THP-1 macrophage (A) proliferation and (B) viability. n=5 per treatment group. *p<0.05, ****p<0.0001.

Results




Figure 3: The effect of oxalate and CFT037 on THP-1 macrophage (A) mitochondrial membrane potential, (B) mitochondrial superoxide, and (C) total reactive oxygen species (ROS) levels. Positive controls: FCCP, LPS, and Menadione. n=5 per treatment group. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.




Figure 4: The effect of oxalate and CFT037 on THP-1 macrophage cellular energetics. Results are presented as mean $\pm$ se; n=5-6 replicates per group.





Figure 5: The effect of oxalate on the anti-bacterial response of THP-1 macrophages over time. *p<0.05, ***p<0.001, ****p<0.0001.

Discussion



- These findings show oxalate impacts macrophage redox status, metabolism, and anti-bacterial response.
- Elucidating the role of CaOx crystals on macrophages during KS disease is critical and should provide new insight to the field since stone formers have increased KS recurrence and associated UTIs.

References

- Tavichakornrakool R, Prasongwattana V, Sungkeeree S et al. Extensive characterizations of bacteria isolated from catheterized urine and stone matrices in patients with nephrolithiasis. *Nephrol. Dial. Transplant. Off. Publ. Eur. Dial. Transpl. Assoc. - Eur. Ren. Assoc.* 2012; 27: 4125-4130
- Barr-Beare E, Saxena V, Hilt EE et al. The Interaction between Enterobacteriaceae and Calcium Oxalate Deposits. *PLoS One* 2015; 10: e0139575.
- Kanlaya R, Naruepantawart O, Thongboonkerd V. Flagellin Is Responsible for Promoting Effects of Viable *Escherichia coli* on Calcium Oxalate Crystallization, Crystal Growth, and Crystal Aggregation. *Frontiers in Microbiology*. 2019;10(2507).
- Borghesi L, Nouvenne A, Meschi T. Nephrolithiasis and urinary tract infections: "the chicken or the egg" dilemma? *Nephrol. Dial. Transplant.* 2012; 27: 3982-3984.
- Jan H, Akbar I, Kamran H et al. Frequency of renal stone disease in patients with urinary tract infection. *J. Ayub Med. Coll. Abbottabad JAMC* 2008; 20: 60-62.
- Umekawa T, Chegwin N, Khan SR. Oxalate ions and calcium oxalate crystals stimulate MCP-1 expression by renal epithelial cells. *Kidney Int* 2002;61(1):105-112.
- Liu Z, Wang T, Yang J, et al. Calcium oxalate monohydrate crystals stimulate monocyte chemoattractant protein-1 and transforming growth factor beta-1 expression in human renal epithelial cells. *Mol Med Rep* 2012;5:1241-1244.

Support & Contact

- NIH grants: K01 DK106284 & R03 DK123542 (TM) and the O'Brien Center P30 DK079337.
- www.ttmitchell-lab.com or email: taneciam@uab.edu

